

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 April 2001 (26.04.2001)

PCT

(10) International Publication Number
WO 01/29007 A1

- (51) International Patent Classification⁷: C07D 231/14, A61K 31/415
- (21) International Application Number: PCT/US00/41239
- (22) International Filing Date: 18 October 2000 (18.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/159,993 18 October 1999 (18.10.1999) US
- (71) Applicant (for all designated States except US): UNIVERSITY OF CONNECTICUT [US/US]; 263 Farmington Avenue, Farmington, CT 06032 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MAKRIYANNIS, Alexandros [US/US]; 348G Foster Drive, Willimantic, CT 06226 (US). LIU, Qian [CN/US]; 1 Northwood Road, Apt. 25, Storrs, CT 06268 (US).
- (74) Agents: YALE, Guy, D. et al.; Alix, Yale & Ristas, LLP, 750 Main Street, Hartford, CT 06103 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- With international search report.
 - Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/29007 A1

(54) Title: PYRAZOLE DERIVATIVES AS CANNABINOID RECEPTOR ANTAGONISTS

(57) Abstract: Novel cannabimimetic pyrazole derivatives are presented which have preferentially high affinities for both of the cannabinoid CB1 or CB2 receptor sites. The improved receptor affinity makes these analogs useful as experimental tools for cannabinoid receptor studies as well as clinically useful agents in individuals and animals for treatment of memory deficits associated with aging or neurological diseases, as anti-obesity agents, as medications for schizophrenia and treatment of septic shock syndrome.

PYRAZOLE DERIVATIVES AS CANNABINOID RECEPTOR ANTAGONISTS

Field of the Invention

The present invention relates generally to pyrazole derivatives and is more particularly concerned with new and improved pyrazole derivatives exhibiting high binding affinities for cannabinoid receptors, pharmaceutical preparations employing these analogs and methods of administering therapeutically effective amounts of the preparations to provide a physiological effect.

5

Background of the Invention

Classical cannabinoids such as the marijuana derived cannabinoid Δ^9 -tetrahydrocannabinol (Δ^9 -THC), as well as endogenous ligands (anandamide) produce their pharmacological effects via their agonist properties at specific cannabinoid receptors in the body. So far, two cannabinoid receptors have been characterized: CB1, a central receptor found in the mammalian brain and peripheral tissues and CB2, a peripheral receptor found only in the peripheral tissues. Compounds that are agonists or antagonists for one or both of these receptors have been shown to provide a variety of pharmacological effects. See, 10 for example, Pertwee, R.G., Pharmacology of cannabinoid CB1 and CB2 receptors, Pharmacol. Ther., (1997) 74:129 - 180 and Di Marzo, V., Melck, D., Bisogno, T., DePetrocellis, L., Endocannabinoids: endogenous cannabinoid receptor ligands with neuromodulatory action, Trends Neurosci. (1998) 21:521 - 528.

20 Over the last few years, a number of potent synthetic cannabinoid agonists have been developed. These agonist materials have helped in the characterization of cannabinoid receptors and with studies of receptor molecular properties.

Cannabinoid antagonists are compounds that bind to one of the CB1 or 25 CB2 receptors but have no effect. There is considerable interest in developing cannabinoid antagonists possessing high affinity for one of the CB1 or CB2 receptors. Such cannabinoid antagonist materials provide a tool to better


understand the mechanisms by which cannabinoid agonists produce their pharmacological effects and for the development of new therapeutic agents.

One class of cannabimimetic antagonists encompasses pyrazole derivatives. Pyrazole analogs have been found to act as antagonists for the CB1 and CB2 receptors, and occasionally to act as agonists for the CB1 and CB2 receptors. Most of the known materials show high receptor affinity for only the CB1 cannabinoid receptor. See for instance, Barth, F. et al, Pyrazole Derivatives, Method Of Preparing Them And Pharmaceutical Compositions In Which They Are Present; U.S. Patent No. 5,624,941 to Barth et al, issued April 29, 1997; Rinaldi-Carmona, M. et al, SR141716A, A Potent And Selective Antagonist Of The Brain Cannabinoid Receptor, *FEBS Lett.* 1994, 350, 240-244; Rinaldi-Carmona, M. et al, Biochemical And Pharmacological Characterization Of SR141716A, The First Potent And Selective Brain Cannabinoid Receptor Antagonist, *Life Sci.* 1995, 56, 1941-1947; and Makriyannis, A., Structure-Activity Relationships Of Pyrazole Derivatives As Cannabinoid Receptor Antagonists, *J. Med. Chem.* 42, 769 - 776, 1999.

Summary of the Invention


The invention includes several novel pyrazole derivatives and physiologically acceptable salts thereof. The invention includes materials selective for either the CB1 or CB2 receptors. Further, some of the analogs have agonistic or antagonistic properties. Pyrazole can be represented by the formula:

25



pyrazole

In one aspect of the invention, modifications were made to the pyrazole structure in the 1, 3, 4 and 5 position of the pyrazole ring. The novel pyrazole derivatives can generally be shown by structural formula 1.



In formula 1, R_1 is a branched or unbranched chain having the structure $(CH_2)_nZ$ where n is an integer from 1 to about 10 and Z is selected from the group consisting of H, halogen, N_3 , NCS (isothiocyanate), CN, OH, OCH_3 , NH_2 and $CH=CH_2$.

R_3 is selected from the group consisting of H and a branched or unbranched chain having the structure $(CH_2)_nCH_3$ where n is an integer from 0 to about 3.

R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 , phenyl and phenyl with at least one substituent from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , and NH_2 .

R_2 is selected from the group consisting of naphthyl,


25

where X is selected from the group consisting of N and CH and Y and Z are each selected from the group consisting of O, N, S and $(CH_2)_n$ where n is an integer from 1 to about 7,

30

where X, Y and Z are each selected from the group consisting of N and CH,

where R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 and phenyl,



where R is selected from the group consisting of H, halogen, N₃, NCS, CN,

5 OH, OCH₃, NH₂ and CH=CH₂.

The novel pyrazole derivatives surprisingly show high binding affinities for either or both of the CB1 and CB2 cannabinoid receptors. Some of the novel pyrazole analogs are cannabinoid receptor antagonists that prevent binding of endogenous agonists to the cannabinoid receptors and thereby block the biological actions of such endogenous agonists. Other novel analogs are cannabinoid receptor agonists. Therefore, the inventive analogs described herein, and physiologically acceptable salts thereof, have high potential when administered in therapeutically effective amounts for providing a physiological effect useful to treat pain, peripheral pain, glaucoma, epilepsy, nausea such as associated with cancer chemotherapy, AIDS Wasting Syndrome, cancer, neurodegenerative diseases including Multiple Sclerosis, Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, mental disorders such as Schizophrenia and depression; to suppress appetite; to reduce fertility; to prevent or reduce diseases associated with motor function such as Tourette's syndrome; to prevent or reduce inflammation; to provide neuroprotection; to modulate the immune system; to produce vasoconstriction or vasodilation and to effect memory enhancement. Thus, another aspect of the invention is the administration of a therapeutically effective amount of an inventive compound, or a physiologically acceptable salt thereof, to an individual or animal to provide a physiological effect.

Additionally, some of the novel pyrazole derivatives have functional moieties such as halogen, azide and isothiocyanate and are potentially useful diagnostic agents *in vivo* (PET, SPECT). Other novel pyrazole derivatives are

radioligands that are potentially useful experimental tools for cannabinoid receptor studies.

Description of Some Preferred Embodiments

5 As used herein a "therapeutically effective amount" of a compound, is the quantity of a compound which, when administered to an individual or animal, results in a sufficiently high level of that compound in the individual or animal to cause a discernible increase or decrease in stimulation of cannabinoid receptors. Physiological effects that result from cannabinoid receptor stimulation include
10 analgesia, decreased nausea resulting from chemotherapy, sedation and increased appetite. Other physiological functions include relieving intraocular pressure in glaucoma patients and suppression of the immune system. Typically, a "therapeutically effective amount" of the compound ranges from about 10 mg/day to about 1,000 mg/day.

15 As used herein, an "individual" refers to a human. An "animal" refers to, for example, veterinary animals, such as dogs, cats, horses and the like, and farm animals, such as cows, pigs and the like.

 The compound of the present invention can be administered by a variety of known methods, including orally, rectally, or by parenteral routes (e.g.,
20 intramuscular, intravenous, subcutaneous, nasal or topical). The form in which the compounds are administered will be determined by the route of administration. Such forms include, but are not limited to, capsular and tablet formulations (for oral and rectal administration), liquid formulations (for oral, intravenous, intramuscular or subcutaneous administration) and slow releasing
25 microcarriers (for rectal, intramuscular or intravenous administration). The formulations can also contain a physiologically acceptable vehicle and optional adjuvants, flavorings, colorants and preservatives. Suitable physiologically to acceptable vehicles may include, for example, saline, sterile water, Ringer's solution, and isotonic sodium chloride solutions. The specific dosage level of
30 active ingredient will depend upon a number of factors, including, for example,

radioligands that are potentially useful experimental tools for cannabinoid receptor studies.

Description of Some Preferred Embodiments

5 As used herein a "therapeutically effective amount" of a compound, is the quantity of a compound which, when administered to an individual or animal, results in a sufficiently high level of that compound in the individual or animal to cause a discernible increase or decrease in stimulation of cannabinoid receptors.


Physiological effects that result from cannabinoid receptor stimulation include
10 analgesia, decreased nausea resulting from chemotherapy, sedation and increased appetite. Other physiological functions include relieving intraocular pressure in glaucoma patients and suppression of the immune system. Typically, a "therapeutically effective amount" of the compound ranges from about 10 mg/day to about 1,000 mg/day.

15 As used herein, an "individual" refers to a human. An "animal" refers to, for example, veterinary animals, such as dogs, cats, horses and the like, and farm animals, such as cows, pigs and the like.

The compound of the present invention can be administered by a variety of known methods, including orally, rectally, or by parenteral routes (e.g.,
20 intramuscular, intravenous, subcutaneous, nasal or topical). The form in which the compounds are administered will be determined by the route of administration. Such forms include, but are not limited to, capsular and tablet formulations (for oral and rectal administration), liquid formulations (for oral, intravenous, intramuscular or subcutaneous administration) and slow releasing
25 microcarriers (for rectal, intramuscular or intravenous administration). The formulations can also contain a physiologically acceptable vehicle and optional adjuvants, flavorings, colorants and preservatives. Suitable physiologically to acceptable vehicles may include, for example, saline, sterile water, Ringer's solution, and isotonic sodium chloride solutions. The specific dosage level of
30 active ingredient will depend upon a number of factors, including, for example,

biological activity of the particular preparation, age, body weight, sex and general health of the individual being treated.

The inventive pyrazole derivatives can generally be described with reference to structural Formula 1:




and physiologically acceptable salts thereof. With reference to structural formula 1, R_1 is a branched or unbranched chain having the structure $(CH_2)_nZ$ where n is an integer from 1 to about 10 and Z is selected from the group consisting of H, halogen, N_3 , NCS, CN, OH, OCH_3 , NH_2 and $CH=CH_2$.

R_3 is selected from the group consisting of H and a branched or unbranched chain having the structure $(CH_2)_nCH_3$ where n is an integer from 0 to about 3.


R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 , phenyl and phenyl with at least one substituent from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , and NH_2 .

R_2 is selected from the group consisting of naphthyl,


30



where X is selected from the group consisting of N and CH and Y and Z are each selected from the group consisting of O, N, S and $(CH_2)_n$ where n is an integer from 1 to about 7,







where X, Y and Z are each selected from the group consisting of N and CH,



5

where R₄, R₅ and R₆ are each selected from the group consisting of halogen, N₃, NCS, OCH₃, CH₃, CH₂CH₃, NO₂, NH₂ and phenyl,



where R is selected from the group consisting of H, halogen, N₃, NCS, CN, OH, OCH₃, NH₂ and CH=CH₂.

10

The following examples are given for purposes of illustration only in order that the present invention may be more fully understood. These examples are not intended to limit in any way the practice of the invention. The above materials were prepared as follows. The prepared cannabimimetic pyrazole derivatives can generally be described with reference to the structures of TABLE 1 below. Naturally, the novel pyrazole derivatives are intended to include physiologically acceptable salts thereof.

15


TABLE 1

	Derivatives	Ki (nM)			Derivatives	Ki (nM)	
		CB ₁	CB ₂			CB ₁	CB ₂
1		5.98	2.51	7		4.73	2.76
2		1.42	0.784	8		1.22	0.615
3		14.6	15.4	9		1.25	0.682
4		7.64	8.46	10		1.34	1.01
5		19.8	6.65	11		3.98	0.965
6		12.2	4.79	12		5.76	0.507

General. Flash column chromatography was carried out using Whatman active silica gel (230 - 400 mesh) and eluents were distilled before use. Solvents for reactions were dried or purified as required. Reactions were carried out under nitrogen atmospheres unless otherwise noted.

General procedure for the preparation of compound 1-5 and 7-8:

Scheme 1:



(a) LiHMDS, ether, then EtO₂CCO₂Et; (b) 5-Chloropentylhydrazine hydrochloride, EtOH; (c) KOH/MeOH; (d) SOCl₂, toluene; (e) Amine, Et₃N, CH₂Cl₂.

5

Lithium salt of ethyl 2,4-dioxo-3-methyl-4-phenylbutanoate. To a magnetically stirred solution of lithium bis(trimethylsilyl)amide (40 ml, 1.0 M solution in hexane, 40 mmol) in diethyl ether (120 mL) was added a solution of propiophenone (5.37 g, 40 mmol) in diethyl ether (50 mL) at

10 78 °C. The mixture was stirred at the same temperature for an additional 45 min, after which diethyl oxalate (6.4 mL, 47 mmol) was added to the mixture. The reaction mixture was allowed to warm to room temperature and stirred for 16 hours (h). The precipitate was filtered, washed with diethyl ether, and dried under vacuum to afford the lithium salt (7.78 g, 83% yield).

15

1-(5-Chloropentyl)-4-methyl-5-phenyl-1H-pyrazole-3-carboxylic acid, Ethyl Ester.

To a magnetically stirred solution of the above lithium salt (2.0 mmol) in 10 mL of ethanol was added a solution of 5-chloropentylhydrazine hydrochloride (2.2 mmol) at room temperature. The resulting mixture was stirred at room temperature for 20 h. The solvent was removed under reduced pressure and the residue was partitioned between ethyl acetate (20 mL) and water (10 mL). The water phase was extracted with ethyl acetate (2x, 15 mL each). The ethyl acetate solution was washed with brine, dried over anhydrous sodium sulfate, filtered, and evaporated. The crude product was purified by flash column


chromatography on silica gel with a petroleum ether/ethyl acetate mixture to afford the ester as a colorless oil.

Compounds 1-5 and 7-8. To a magnetically stirred solution of the above ethyl ester (3.4 mmol) in methanol (15 mL) was added a solution of potassium hydroxide (8.6 mmol) in methanol (12 mL). The mixture was heated under reflux for 3 hours. The cooling reaction mixture was then poured into 10 mL of water and acidified with 10% hydrochloric acid. The precipitate was filtered, washed with water, and dried under vacuum to yield the corresponding acid (1.4g, 100% yield) as a white solid.

A solution of the crude acid (3.4 mmol) and thionyl chloride (10.3 mmol) in toluene (15 mL) was refluxed for 3 hours. The solvent was evaporated under reduced pressure. The residue was then redissolved in 40 mL of toluene and evaporated to yield the crude carboxylic chloride as an oil.

A solution of the carboxylic chloride (31.5 mmol) in dichloromethane (160 mL) was added dropwise to a solution of an appropriate amine (47.2 mmol) and triethylamine (6.5 mL, 46.7 mmol) in dichloromethane (90 mL) at 0°C. After stirring at room temperature for 3 h, brine was added to the reaction mixture, which was extracted with dichloromethane (3x, 200 mL each). The combined extracts were washed with brine, dried over anhydrous sodium sulfate, filtered, and evaporated. Flash column chromatography on silica gel with a petroleum ether/acetone (4:1) mixture gave carboxamide 1-5 and 7-8.

Scheme 2:




(a) PCC, CH₂Cl₂; (b) t-Butyl carbazate, hexane; (c) BH₃/THF, THF; (d) 6N HCl.

5-Chloropentylhydrazine hydrochloride. A hexane solution of 5-chloropentylaldehyde (10.0 mmol) and *tert*-butyl carbazate (1.32 g, 10.0 mmol) was refluxed for 20 min. After cooling to the room temperature, the crystallized *tert*-butyl carbazate derivative was collected by filtration and dried in vacuum.

- 5 A 1.0 M solution of borane tetrahydrofuran complex in tetrahydrofuran (10.0 mL, 10.0 mmol) was added to the solid *tert*-butyl carbazate derivative (10.0 mmol), the resulting mixture was allowed to stir at room temperature for 10 min, and then 6N hydrochloric acid (5.0 mL) was added dropwise. The reaction mixture was refluxed for 10 min and evaporated to dryness under reduced
- 10 pressure. Tetrahydrofuran was added to the residue, after which boric acid was removed by filtration. After removal of the solvent under reduced pressure, the residue was crystallized from a solution of tetrahydrofuran and diethyl ether to give 5-chloropentylhydrazine as its hydrochloride salt (71 % yield).

15 **Scheme 3:**



Preparation of compound 9:


- To a magnetically stirred solution of compound 2 (0.40 g, 0.91 mmol) in
- 20 acetonitrile (15 mL) was added a 1.0 M solution of tetrabutylammonium fluoride (4.5 mL, 4.5 mmol) in tetrahydrofuran and the mixture was refluxed overnight. The reaction mixture was then quenched by saturated aqueous ammonium

chloride and extracted with diethyl ether (3×, 50 mL each). The combined extracts were washed with brine, dried over anhydrous sodium sulfate, filtered and evaporated. Purification by flash column chromatography on silica gel with a petroleum ether/acetone (9:1) mixture gave compound 9 as a white solid (0.296 g, 77% yield).

Preparation of compound 10:

To a magnetically stirred solution of compound 2 (1.12 g, 2.6 mmol) in acetone (25 mL) was added sodium iodide (1.72g, 11.5 mmol). The reaction mixture was refluxed for 25 hours and evaporated to dryness under reduced pressure. The residue was partitioned between diethyl ether (100 mL) and water (40 mL), and the water phase was extracted with diethyl ether (3×, 30 mL each). The combined extracts were washed with brine, dried over anhydrous sodium sulfate, filtered, and evaporated. Purification by flash column chromatography on silica gel with a methylene chloride/acetone (30:1) mixture gave compound 10 as a white solid (1.27 g, 94 % yield).

Scheme 4:



(a) NaN_3 , DMF; (b) CS_2 , Ph_3P , THF.

Preparation of compound 11:

To a magnetically stirred solution of compound 10 (0.675 g, 1.3 mmol) in anhydrous N,N-dimethylformamide (17 mL) was added sodium azide (0.83 g,

12.7 mmol). The resulting mixture was stirred at room temperature for 40 hours. Brine was then added and the reaction mixture was extracted with diethyl ether (3×, 20 mL each). The combined extracts were washed with brine, dried over anhydrous sodium sulfate, filtered and evaporated. The residue was
5 purified by flash column chromatography on silica gel with a methylene dichloride/acetone (50:1) mixture to afford compound 11 as a white solid (0.203 g, 35.8 % yield).

Preparation of compound 12:

10 To a magnetically stirred solution of compound 11 (0.359 g, 0.8 mmol) in


12.7 mmol). The resulting mixture was stirred at room temperature for 40 hours. Brine was then added and the reaction mixture was extracted with diethyl ether (3x, 20 mL each). The combined extracts were washed with brine, dried over anhydrous sodium sulfate, filtered and evaporated. The residue was
 5 purified by flash column chromatography on silica gel with a methylene dichloride/acetone (50:1) mixture to afford compound 11 as a white solid (0.203 g, 35.8 % yield).

Preparation of compound 12:

10 To a magnetically stirred solution of compound 11 (0.359 g, 0.8 mmol) in tetrahydrofuran (5 mL) was added triphenylphosphine (0.32 g, 1.22 mmol), followed by carbon disulfide (1.44 mL, 24 mmol). The reaction mixture was stirred at room temperature for 70 hours and then evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel
 15 with methylene dichloride to afford compound 12 (0.288 g, 77.6% yield).

Preparation of compound 6:

Scheme 5:



20 (a) NaOCH₃/CH₃OH, EtO₂CCO₂Et; (b) 5-Chloropentylhydrazine hydrochloride, HAc;
 (c) KOH/MeOH; (d) SOCl₂, toluene; (e) Amine, Et₃N, CH₂Cl₂.

Sodium salt of methyl benzoylpyruvate. 1.3 g of sodium was dissolved in 25 mL of anhydrous methanol. A mixture of 5.8 mL of acetophenone and 6.7 mL of
 25 diethyl oxalate in 60 mL of methanol was then added, the temperature being

kept below 10 °C. The reaction mixture was then stirred at room temperature for 3 hours, after which 100 mL of dry ether was added. Stirring was continued for 20 min, the mixture was filtered and precipitate was washed with ether and dried under vacuum to give 6.32 g of the expected sodium salt.

5

1-(5-Chloropentyl)-5-phenyl-1*H*-pyrazole-3-carboxylic acid, Methyl Ester. A suspension of 0.605 g of the sodium salt obtained above and 0.502 g of 5-chloropentylhydrazine hydrochloride in 6.5 mL of acetic acid was refluxed for 4 hours. After cooling, the mixture was poured on to 6.5 g of ice and the crystals
10 obtained were filtered off, washed with water and dried under vacuum to give 0.42 g of ester.

Compound 6. Compound 6 was prepared from the methyl ester according to the procedure described for the compound 1-5 and 7-8.

15

The materials were tested for CB2 receptor binding affinity and for CB1 receptor affinity (to determine selectivity for the CB2 receptor). As used herein, "binding affinity" is represented by the IC_{50} value which is the concentration of an analog required to occupy 50% of the total number (B_{max}) of the receptors.

The lower the IC_{50} value, the higher the binding affinity. As used herein an
20 analog is said to have "binding selectivity" if it has higher binding affinity for one receptor compared to the other receptor; e.g. a cannabinoid analog which has an IC_{50} of 0.1 nM for CB1 and 10 nM for CB2, is 100 times more selective for the CB1 receptor. The binding affinities (K_i) are expressed in nanomoles (nM) and are listed in TABLE 1.

25

For the CB1 receptor binding studies, membranes were prepared from rat forebrain membranes according to the procedure of P.R. Dodd et al, A Rapid Method for Preparing Synaptosomes: Comparison with Alternative Procedures, Brain Res., 107 - 118 (1981). The binding of the novel analogues to the CB1 cannabinoid receptor was assessed as described in W.A. Devane et al,
30 Determination and Characterization of a Cannabinoid Receptor in a Rat Brain,

Mol. Pharmacol., 34, 605 - 613 (1988) and A. Charalambous et al, 5'-azido Δ^8 -THC: A Novel Photoaffinity Label for the Cannabinoid Receptor, J. Med. Chem., 35, 3076 - 3079 (1992) with the following changes. The above articles are incorporated by reference herein.

5 Membranes, previously frozen at -80°C , were thawed on ice. To the stirred suspension was added three volumes of TME (25mM Tris-HCl buffer, 5 mM MgCl_2 and 1 mM EDTA) at pH 7.4. The suspension was incubated at 4°C for 30 min. At the end of the incubation, the membranes were pelleted and washed three times with TME.

10 The treated membranes were subsequently used in the binding assay described below. Approximately 30 μg of membranes were incubated in silanized 96-well microtiter plate with TME containing 0.1% essentially fatty acid-free bovine serum albumin (BSA), 0.8 nM $[^3\text{H}]$ CP-55,940, and various concentrations of test materials at 30°C for 1 hour. The samples were filtered
15 using Packard Filtermate 196 and Whatman GF/C filterplates and washed with wash buffer (TME containing 0.5% BSA). Radioactivity was detected using MicroScint 20 scintillation cocktail added directly to the dried filterplates, and the filterplates were counted using a Packard Instruments Top-Count. Nonspecific binding was assessed using 100 nM CP-55,940. Data collected from three
20 independent experiments performed with duplicate determinations was normalized between 100% and 0% specific binding for $[^3\text{H}]$ CP-55,940, determined using buffer and 100 nM CP-55,940. The normalized data was analyzed using a 4-parameter nonlinear logistic equation to yield IC_{50} values. Data from at least two independent experiments performed in duplicate was used
25 to calculate IC_{50} values which were converted to K_i values using the assumptions of Cheng et al, Relationship Between the Inhibition Constant (K_i) and the concentration of Inhibitor which causes 50% Inhibition (IC_{50}) of an Enzymatic Reaction, Biochem. Pharmacol., 22, 3099-3102, (1973), which is incorporated by reference herein.

For the CB2 receptor binding studies, membranes were prepared from frozen mouse spleen essentially according to the procedure of P.R. Dodd et al, A Rapid Method for Preparing Synaptosomes: Comparison with Alternative Procedures, Brain Res., 226, 107 - 118 (1981) which is incorporated by
5 reference herein. Silanized centrifuge tubes were used throughout to minimize receptor loss due to adsorption. The CB2 binding assay was conducted in the same manner as for the CB1 binding assay. The binding affinities (K_i) were also expressed in nanomoles (nM).

Compound SR141716A, a known pyrazole derivative, has a cannabinoid
10 receptor affinity (K_i) of 11.5 nM for the CB1 receptor and 1640 nM for the CB2 receptor. As can be seen from the results in TABLE 1, all of the inventive compounds have receptor affinities much higher than compound SR141716A for at least one of the CB1 or CB2 receptors. In fact, most of the inventive pyrazole derivatives have receptor affinities much higher (lower numerically) than
15 compound SR141716A for both of the CB1 and CB2 receptors.

The physiological and therapeutic advantages of the inventive materials can be seen from the above disclosure and also with additional reference to the following references, the disclosures of which are hereby incorporated by reference. Arnone M., Maruani J., Chaperon P, *et al*, Selective inhibition of
20 sucrose and ethanol intake by SR141716, an antagonist of central cannabinoid (CB1) receptors, Psychopharmacol, (1997) 132, 104-106. Colombo G, Agabio R, Diaz G. et al: Appetite suppression and weight loss after the cannabinoid antagonist SR141716. Life Sci. (1998) 63-PL13-PL117. Simiand J, Keane M, Keane PE, Soubrie P: SR 141716, A CB1 cannabinoid receptor antagonist,
25 selectively reduces sweet food intake in marmoset. Behav. Pharmacol (1998) 9:179-181. Brotchie JM: Adjuncts to dopamine replacement a pragmatic approach to reducing the problem of dyskinesia in Parkinson's disease. Mov. Disord. (1998) 13:871-876. Terranova J-P, Storme J-J Lafon N et al: Improvement of memory in rodents by the selective CB1 cannabinoid receptor
30 antagonist, SR 141716. Psycho-pharmacol (1996) 126:165-172. Hampson AL

- Grimaldi M, Axirod J, Wink D: Cannabidiol and (-) Δ^9 tetrahydrocannabinol are neuroprotective antioxidants. Proc. Natl Acad Sci. USA (1998) 95:8268-8273.
- Buckley NE, McCoy KI, Mpzey E et al Immunomodulation by cannabinoids is absent in mice deficient for the cannabinoid CB₂ receptor. Eur. J Pharmacol
- 5 (2000) 396:141-149. Morgan Dr: Therapeutic Uses of Cannabis. Harwood Academic Publishers, Amsterdam. (1997). Joy JE, Wagtson SJ, Benson JA: Marijuana and Medicine Assessing the Science Base. National Academy Press, Washington, DC, USA (1999). Shen M, Thayer SA: Cannabinoid receptor agonists protect cultured rat hippocampal neurons from excitotoxicity. Mol.
- 10 Pharmacol (1996) 54:459-462. DePetrocellis L, Melck D, Palmisano A. et al: The endogenous cannabinoid anandamide inhibits human breast cancer cell proliferation. Proc Natl. Acad. Sci USA (1998) 95:8375-8380. Green K. Marijuana smoking vs. cannabinoids for glaucoma therapy. Arch. Ophthalmol. (1998) feb 433-437. Hemming M, Yellowlees PM, Effective treatment of
- 15 Tourette's syndrome with marijuana. J. Psychopharmacol, (1993) 7:389-391. Muller-Vahl KB, Schneider U, Kolbe H, Emrich, HM. Treatment of Tourette's syndrome with delta-9-tetrahydrocannabinol. Am. J. Psychiat. (1999) 156-195. Muller-Vahl KB, Kolbe H, Schneider U, Emrich, HM Cannabis in movement disorders. Porsch. Kompimentarmed (1999) 6 (suppl. 3) 23-27. Consroe P,
- 20 Musty R, Rein J, Tillery W, Pertwee R. The perceived effects of smoked cannabis on patients with multiple sclerosis, Eur. Neurol. (1997) 38-44-48. Pinnegan-Ling D, Musty R. Marinol and phantom limb pain: a case study. Proc Inv. Cannabinoid Rea. Sec. (1994):53. Brenneisen R, Pgli A, Elsohly MA, Henn V. Spiess Y: The effect of orally and rectally administered Δ^9 -tetrahydrocannabinol on spasticity,
- 25 a pilot study with 2 patients. Int. J. Clin Pharmacol Ther. (1996) 34:446-452. Martyn CN, Illis LS, Thom J. Nabilone in the treatment of multiple sclerosis. Lancet (1995) 345:579. Maurer M, Henn V, Dittrich A, Hofmann A. Delta-9-tetrahydrocannabinol shows antispastic and analgesic effects in a single case double-blind trial. Eur. Arch. Psychiat. Clin. Neurosci. (1990), 240:1-4. Herzberg
- 30 U, Eliav E, Bennett GJ, Kopin IJ: The analgesic effects of R(+) WIN 55,212-2


- mesylate, a high affinity cannabinoid agonist in a rare model of neuropathic pain. Neurosci. Letts. (1997) 221:157-160. Richardson JD, Kilo S. Hargreaves KM, Cannabinoids reduce dryperalgesia and inflammation via interaction with peripheral CB1 receptors. Pain (1998) 75:111-119. Ricardson JD, Aanonsen I, 5 Hargreaves KM: Antihyperalgesic effects of a spinal cannabinoids. Eur. J. Pharmacol. (1998) 346:145-153. Calignano A, La Rana G. Diuffrida A, Piomelli D: Control of pain initiation by endogenous cannabinoids. Nature (1998) 394:277-291. Wagner JA, Varga K, Jarai Z, Kunos G: Mesenteric vasodilation mediated by endothelia anandamide receptors. Hypertension (1999) 33:429- 10 434. Schuel, H., Burkman, L.J., Picone, R.P., Bo, T., Makriyannis, A., Cannabinoid receptors in human sperm. Mol. Biol. Cell., (1997) (8), 325a.

The inventive analogs described herein, and physiologically acceptable salts thereof, have high potential when administered in therapeutically effective amounts for providing a physiological effect useful to treat pain, peripheral pain, 15 glaucoma, epilepsy, nausea such as associated with cancer chemotherapy, AIDS Wasting Syndrome, cancer, neurodegenerative diseases including Multiple Sclerosis, Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, mental disorders such as Schizophrenia and depression; to suppress appetite; to reduce fertility; to prevent or reduce diseases associated with motor function 20 such as Tourette's syndrome; to prevent or reduce inflammation; to provide neuroprotection; to modulate the immune system; to produce vasoconstriction or vasodilation and to effect memory enhancement. Thus, another aspect of the invention is the administration of a therapeutically effective amount of an inventive compound, or a physiologically acceptable salt thereof, to an individual 25 or animal to provide a physiological effect.

Those skilled in the art will recognize, or be able to ascertain with no more than routine experimentation, many equivalents to the specific embodiments of the invention disclosed herein. Such equivalents are intended to be encompassed by the scope of the invention.

What Is Claimed Is:

1. A compound of the formula




or a physiologically acceptable salt thereof, wherein:

R_1 is a branched or unbranched chain having the structure $(CH_2)_nZ$ where n is an integer from 1 to about 10 and Z is selected from the group consisting of H, halogen, N_3 , NCS, CN, OH, OCH_3 , NH_2 and $CH=CH_2$;


R_3 is selected from the group consisting of H and a branched or unbranched chain having the structure $(CH_2)_nCH_3$ where n is an integer from 0 to about 3;

R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 , phenyl and phenyl with at least one substituent from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , and NH_2 ; and


R_2 is selected from the group consisting of naphthyl,






where X is selected from the group consisting of N and CH and Y and Z are each selected from the group consisting of O, N, S and $(CH_2)_n$ where n is an integer from 1 to about 7,



where X , Y and Z are each selected from the group consisting of N and CH,




where R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 and phenyl,



where R is selected from the group consisting of H, halogen, N_3 , NCS, CN, OH, OCH_3 , NH_2 and $CH=CH_2$.

2. A method of preferentially binding to the cannabinoid receptors in an individual or animal comprising administering to the individual or animal a therapeutically effective amount of a compound having the formula



and physiologically acceptable salts thereof, wherein:


R_1 is a branched or unbranched chain having the structure $(CH_2)_nZ$ where n is an integer from 1 to about 10 and Z is selected from the group consisting of H, halogen, N_3 , NCS, CN, OH, OCH_3 , NH_2 and $CH=CH_2$;

R_3 is selected from the group consisting of H and a branched or unbranched chain having the structure $(CH_2)_nCH_3$ where n is an integer from 0 to about 3;


R_4 , R_5 and R_6 are each selected from the group consisting of halogen,

N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 , phenyl and phenyl with at least one substituent from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , and NH_2 ; and


R_2 is selected from the group consisting of naphthyl,






where X is selected from the group consisting of N and CH and Y and Z are each selected from the group consisting of O, N, S and $(CH_2)_n$ where n is an integer from 1 to about 7,



where X, Y and Z are each selected from the group consisting of N and CH,




where R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 , phenyl and



where R is selected from the group consisting of H, halogen, N_3 , NCS, CN, OH, OCH_3 , NH_2 and $CH=CH_2$.

3. A pharmaceutical composition containing a therapeutically effective amount of a compound having the formula




and physiologically acceptable salts thereof, wherein:

R_1 is a branched or unbranched chain having the structure $(CH_2)_nZ$ where n is an integer from 1 to about 10 and Z is selected from the group consisting of H, halogen, N_3 , NCS, CN, OH, OCH_3 , NH_2 and $CH=CH_2$;


R_3 is selected from the group consisting of H and a branched or unbranched chain having the structure $(CH_2)_nCH_3$ where n is an integer from 0 to about 3;

R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 , phenyl and phenyl with at least one substituent from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , and NH_2 ; and


R_2 is selected from the group consisting of naphthyl,







where X is selected from the group consisting of N and Ch and Y and Z are each selected from the group consisting of O, N, S and $(CH_2)_n$ where n is an integer from 1 to about 7,



where X , Y and Z are each selected from the group consisting of N and CH,



where R_4 , R_5 and R_6 are each selected from the group consisting of halogen, N_3 , NCS, OCH_3 , CH_3 , CH_2CH_3 , NO_2 , NH_2 , phenyl and



where R is selected from the group consisting of H, halogen, N_3 , NCS, CN, OH, OCH_3 , NH_2 and $CH=CH_2$.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/41239

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 231/14; A61K 31/415

US CL : 548/374.1; 514/406

According to international Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 548/374.1; 514/406

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	US 6,127,399 A (YUAN) 03 October 2000(03/10/00), see entire document, especially claims 1-4.	1-3

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claims, or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
I document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 JANUARY 2001

Date of mailing of the international search report

13 MAR 2001

Name and mailing address of the ISA-US
Commissioner of Patents and Trademarks
Box PCT
Washington, D. 20231

Facsimile No. (703) 305-3230

Authorized officer

R. W. RAMSUER

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/41239

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-3.(parts)
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Please See Extra Sheet.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/41239

BOX I. OBSERVATIONS WHERE CLAIMS WERE FOUND UNSEARCHABLE

2. Where no meaningful search could be carried out, specifically:

The multitude of variables and their permutations and combinations (e.g. R1, R2, R3, R4, X, Y, Z, etc.) result in claimed subject matter which is so voluminous in scope that no meaningful complete search can be given. Note also that the claimed subject matter lacks a significant structural element that qualifies as the special technical feature that clearly defines a contribution over the art. The subject matter claimed contains a phenylpyrazolecarbonyl N group which does not define a contribution over the prior art. Therefore, the first discernible invention as found in the Examples (i.e. compounds 1-2 and 4-12, the pharmaceutical compositions and the method of preferentially binding to the cannabinoid receptors utilizing said compounds) as covered by claims 1-3, has been searched.